

Connexin 36 antibody, Internal

Cat. No. GTX47483

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC
Reactivity	Human, Mouse

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 ug/ml
IHC	Assay dependent

Not tested in other applications.

Calculated MW 36 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to an Internal region of Human Connexin 36.
Purification	Affinity Purified
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX47483 IHC Image

IHC analysis of mouse brain tissue using GTX47483 GJD2 antibody at 1 μ g/ml. Left to right : DAPI, GJD2, parvalbumin, merge.


GTX47483 WB Image

WB analysis of 293T cells using GTX47483 GJD2 antibody at 1 μ g/ml.


GTX47483 WB Image

WB analysis of Jurkat cells using GTX47483 GJD2 antibody at 0.2-1 μ g/ml.



For full product information, images and publications, please visit our [website](https://www.genetex.com).