

Aquaporin 12 antibody (FITC)

Cat. No. GTX47924

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IP, ELISA, IHC
Reactivity	Mouse, Rat

Package 50 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IP	Assay dependent
ELISA	Assay dependent
IHC	Assay dependent
Not tested in other applications.	

Calculated MW 31 kDa. (<u>Note</u>)

Form Liquid Buffer Tris/Glycine, 0.5% BSA, 30% Glycerol
Buffer Tris/Glycine, 0.5% BSA, 30% Glycerol
Preservative 0.02% Sodium azide
Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration 0.61-0.67 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen Synthetic peptide taken within amino acid region 240-290 on mouse Aquaporin 12 protein.
Purification Purified IgG
Conjugation Fluorescein isothiocyanate (FITC) Wavelength



For full product information, images and publications, please visit our website.

Date 2025 / 11 / 04 Page 1 of 2



For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 04 Page 2 of 2