

Aquaporin 10 antibody (FITC)

Cat. No. GTX47928

| | |
|---------------------|----------------------------|
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, ICC/IF, IP, ELISA, IHC |
| Reactivity | Human, Mouse, Rat |

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | Assay dependent |
| ICC/IF | Assay dependent |
| IP | Assay dependent |
| ELISA | Assay dependent |
| IHC | Assay dependent |

Not tested in other applications.

Calculated MW 32 kDa. ([Note](#))

Properties

| | |
|----------------------|--|
| Form | Liquid |
| Buffer | Tris/Glycine, 0.5% BSA, 30% Glycerol |
| Preservative | 0.02% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light. |
| Concentration | 0.64-0.88 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Synthetic peptide taken within amino acid region 250-300 on human Aquaporin 10 protein. |
| Purification | Purified IgG |
| Conjugation | Fluorescein isothiocyanate (FITC) Wavelength |



For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).