

Human TGF Beta 1 protein (active)

Cat. No. GTX48131-pro

Applications	Functional Assay
Species	Human

Package
10 µg

Applications

Application Note

The ED₅₀ as determined by the ability of TGF-β2 to inhibit the mouse IL-4-dependent proliferation of mouse HT-2 cells is <0.2 ng/ml, corresponding to a specific activity of > 5 x 10⁶ units/mg.

Properties

Form	Lyophilized powder
Buffer	Batch dependent (please contact us for details)
Storage	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
Region/Sequence	ALDTNYCFSS TEKNCCVRQL YIDFRKDLGW KWIHEPKGYH ANFLGPCPY IWSLDTQYSK VLALYNQHNP GASAAPCCVP QALEPLPIVY YVGRKPKVEQ LSNMIVRSCK CS
Expression System	CHO cells
Purity	> 98% by SDS-PAGE and HPLC.
Endotoxin	< 1 EU/µg
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).