

Human FGF8 protein (active)

Cat. No. GTX48450-pro

Applications	Functional Assay	Package
Species	Human	25 µg

Applications

Application Note

The ED₅₀ was determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF-receptors is ≤ 0.5 ng/ml, corresponding to a specific activity of $\geq 2 \times 10^6$ units/mg.

Properties

Form	Lyophilized powder
Buffer	Batch dependent (please contact us for details)
Storage	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
Region/Sequence	MQTVQSSPN FTQHVREQSL VTDQLSRRLI RTYQLYSRTS GKHVQVLANK RINAMAEDGD PFAKLIVETD TFGSRVRVRG AETGLYICMN KKGKLIAKSN GKGKDCVFTE IVLENNYTAL QNAKYEGWYM AFTRKGRPRK GSCTRQHQRE VHFMKRLPRG HHTTEQSLRF EFLNYPPFTR SLRGSQRTWA PEPR
Expression System	E. coli
Purity	> 95% by SDS-PAGE and HPLC.
Endotoxin	< 1 EU/µg
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 1 of 1