

## Human FGF2 protein (active)

Cat. No. GTX48452-pro

Application	Functional Assay
Species	Human

Package  
50 µg

## APPLICATION

## Application Note

Assay #1: The ED<sub>50</sub> was determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF receptors is  $\leq 0.5$  ng/ml, corresponding to a specific activity of  $\geq 2 \times 10^6$  units/mg. Assay #2: The ED<sub>50</sub> was determined by a cell proliferation assay using balb/c 3T3 cells is  $\leq 0.1$  ng/ml, corresponding to a specific activity of  $\geq 1 \times 10^7$  units/mg.

## PROPERTIES

Form	Lyophilized powder
Buffer	Batch dependent (please contact us for details)
Storage	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
Region/Sequence	AAGSITTLPA LPEDGGSGAF PPGHFKDPKR LYCKNGGFFL RIHPDGRVDG VREKSDPHIK LQLQAEERGV VSIKGVCANR YLAMKEDGRL LASKCVTDEC FFERLESNN YNTYRSRKYT SWYVALKRTG QYKLGSKTGP GQKAILFLPM SAKS
Expression System	E. coli
Purity	> 95% by SDS-PAGE and HPLC.
Endotoxin	< 1 EU/µg
Conjugation	Unconjugated

## Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).