

LDB1 antibody

Cat. No. GTX48750

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ELISA, ChIP assay
Reactivity	Human

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ELISA	1:425000
ChIP assay	Assay dependent

Not tested in other applications.

Calculated MW 47 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.8 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide corresponding to a C-Terminal region of mouse LDB1 protein.
Purification	Purified by antigen-affinity chromatography. From serum
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTx48750 WB Image

Western blot using GeneTex's affinity purified anti-LDB1 antibody shows detection of LDB1 protein (arrowhead) in Jurkat whole cell lysate. Approximately 30 µg of lysate was loaded prior to separation and transfer to nitrocellulose. Primary antibody was used at a 1:1,800 dilution in 5% BLOTTO in PBS reacted overnight at 4°C. The membrane was washed and reacted with a 1:20,000 dilution of DyLight800 conjugated goat anti-Rabbit IgG [H&L] MX for 45 min at room temperature (800 nm channel, green). Molecular weight estimation was made by comparison to prestained MW markers in lane M (700 nm channel, red). Fluorescence image was captured using the OdysseyR Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc. Other detection systems will yield similar results.



GTx48750 WB Image

WB analysis of Jurkat whole cell lysate using GTx48750 LDB1 antibody.

Loading : 30 µg

Dilution : 1:800



For full product information, images and publications, please visit our [website](https://www.genetex.com).