

CRASP-2 antibody

Cat. No. GTX48800

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ELISA
Reactivity	Borrelia burgdorferi

Package
50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000
ELISA	1:2000

Not tested in other applications.

Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	MBP-fusion protein corresponding to Borrelia burgdorferi CRASP-2 protein.
Purification	Protein A purified From serum
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTx48800 WB Image

WB analysis of various samples using GTx48800 CRASP-2 antibody.


Lane 1 : Protein ladder

Lane 2 : MBP-CRASP-2 fusion protein

Lane 3 : MBP

Loading : 0.1 µg

Dilution : 1:1000



GTx48800 WB Image

Western Blot showing detection of 0.1 µg of recombinant CRASP-2 protein. Lane 1: Molecular weight markers. Lane 2: MBP-CRASP-2 fusion protein (arrow; expected MW = 67.8 kDa). Lane 3: MBP alone. Protein was run on a 4-20% gel, then transferred to 0.45 µm nitrocellulose. After blocking with 1% BSA-TTBS overnight at 4°C, primary antibody was used at 1:1000 at room temperature for 30 min. HRP-conjugated Goat-Anti-Rabbit secondary antibody was used at 1:40,000 in blocking buffer and imaged on the VersaDoc MP 4000 imaging system (Bio-Rad).



For full product information, images and publications, please visit our [website](https://www.genetex.com).