

Borrelia burgdorferi Osp A antibody

Cat. No. GTX48804

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ELISA
Reactivity	Borrelia burgdorferi

Package

50 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000
ELISA	>1:5000

Not tested in other applications.

Calculated MW 29 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.12 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	MBP-fusion protein corresponding to Borrelia burgdorferi OspA protein.
Purification	Protein A purified From serum
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


Date 2026 / 01 / 11 Page 1 of 2

DATA IMAGES




GTX48804 WB Image

Western blot showing detection of 0.1 µg of recombinant OspA protein. Lane 1: Molecular weight markers. Lane 2: MBP-OspA fusion protein (arrow; expected MW: 70.5 kDa). Lane 3: MBP alone. Protein was run on a 4-20% gel, then transferred to 0.45 µm nitrocellulose. After blocking with 1% BSA-TTBS overnight at 4°C, primary antibody was used at 1:1000 at room temperature for 30 min. HRP-conjugated Goat-Anti-Rabbit secondary antibody was used at 1:40,000 in blocking buffer and imaged on the VersaDoc MP 4000 imaging system (Bio-Rad).



GTX48804 WB Image

WB analysis of various samples using GTX48804 Borrelia burgdorferi Osp A antibody.
Lane 1 : Protein ladder
Lane 2 : MBP-Osp A fusion protein
Lane 3 : MBP
Loading : 0.1 µg
Dilution : 1:1000



GTX48804 WB Image

WB analysis of various samples using GTX48804 Borrelia burgdorferi Osp A antibody.
Lane 1 : Protein ladder
Lane 2 : 50 ng Osp A
Lane 3 : 20 ng Osp A
Lane 4 : 10 ng Osp A
Lane 5 : Blank
Lane 6 : 50 ng Osp B
Lane 7 : 50 ng Osp C
Dilution : 1:1000



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 11 Page 2 of 2