

## MED25 antibody, N-term

## Cat. No. GTX49103

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ChIP assay
Reactivity	Human

## Package

50 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 µg/ml
ChIP assay	Assay dependent

Not tested in other applications.

Calculated MW 78 kDa. ([Note](#))

## Properties


Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to a N-terminal region of Human MED25
Purification	Affinity Purified
Conjugation	Unconjugated
	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 1 of 2

## DATA IMAGES




## GTX49103 ChIP assay Image

ChIP analysis of HCT116 cells using GTX49103 MED25 antibody. Use 0.5 $\mu$ g antibody GTX49103 MED25 antibody for ChIP assay. Quiescent human colon carcinoma HCT116 cultures were treated with 10% FBS for three time points (0, 15, 30min) or (0, 30, 60min) were used in Matrix-ChIP and real-time PCR assays at EGR1 gene (Exon1) and 15kb upstream site.

(+) : Add blocking peptide

(-) : no blocking peptide



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 2 of 2