

c-Met (phospho Tyr1349) antibody

Cat. No. GTX50270

Host	Rabbit
Clonality	Polyclonal
Isotype	lgG
Applications	WB
Reactivity	Human

Package 100 μΙ

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:1000
Not tested in other applications.	

Calculated MW 156 kDa. (Note)


Properties	
Form	Liquid
Buffer	PBS, 150mM NaCl, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide sequence around phosphorylation site of tyrosine 1349 (E-H-Y(p)-V-H) derived from human c-Met.
Purification	Purified by antigen-affinity chromatography. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.

Date 2025 / 06 / 18 Page 1 of 2

DATA IMAGES



GTX50270 WB Image

WB analysis of extracts from HepG2 cells using GTX50270 c-Met (phospho Tyr1349) antibody. Lane 1: Primary antibody pre-incubated with the antigen specific peptide



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 18 Page 2 of 2