

EGFR (phospho Tyr1086) antibody

Cat. No. GTX50289

Host	Rabbit	Package
Clonality	Polyclonal	100 μ l
Isotype	IgG	
Applications	WB	
Reactivity	Human	

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:1000
Not tested in other applications.	
Calculated MW	134 kDa. (Note)
Product Note	The full-length EGFR includes the 24 aa signal peptide that is cleaved during EGFR maturation. The Tyr1086 of mature EGFR is equivalent to Tyr1110 of uncleaved EGFR.

Properties


Form	Liquid
Buffer	PBS, 150mM NaCl, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide sequence around phosphorylation site of tyrosine 1086 (P-V-Y(p)-H-N) derived from human EGFR.
Purification	Purified by antigen-affinity chromatography. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 01 Page 1 of 2

DATA IMAGES



GTX50289 WB Image

WB analysis of extracts from HUVEC cells untreated or treated with EGF using GTX50289 EGFR (phospho Tyr1086) antibody.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 01 Page 2 of 2