

## c-Myc antibody

## Cat. No. GTX50395

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package  
100 µl

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:1000
IHC-P	1:50-1:100

Not tested in other applications.

Calculated MW 49 kDa. ([Note](#))

## Properties


Form	Liquid
Buffer	PBS, 150mM NaCl, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide sequence around aa.356~360 (R-R-T-H-N) derived from human c-Myc.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 10 Page 1 of 2

## DATA IMAGES

**GTX50395 WB Image**

WB analysis of extracts from HL-60, MCF, and K562 cells using GTX50395 c-Myc antibody.

**GTX50395 IHC-P Image**

IHC-P analysis of human breast carcinoma tissue using GTX50395 c-Myc antibody.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 10 Page 2 of 2