

CD73 antibody [4G4]

Cat. No. GTX54475

| Host | Mouse |
|--------------|--|
| Clonality | Monoclonal |
| Isotype | lgG1 |
| Applications | WB, ICC/IF, IHC-Fr, FCM, IP, Neutralizing/Inhibition |
| Reactivity | Human |

Package 50 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|-------------------------|----------------------|
| WB | Assay dependent |
| ICC/IF | Assay dependent |
| IHC-Fr | Assay dependent |
| FCM | Assay dependent |
| IP | Assay dependent |
| Neutralizing/Inhibition | Assay dependent |

Note: HUVEC cells were seeded on gelatin-coated coverslips and stained with antibody.

Tissue sections were fixed in acetone.- As positive control anti-CD31 was used and as negative control an isotype antibody.

Antibody 4G4 functions as an inhibitor of lymphocyte binding to HUVEC. The antibody was functionally tested by incubating the lymphocytes with the antibody before adding the lymphocytes to an EC monolayer. Furthermore the monoclonal antibody 4G4 causes a reduction in CD73 expression on lymphocytes and reduces enzyme activity.

Not tested in other applications.

| Calculated MW | 63 kDa. (<u>Note</u>) |
|---------------|--|
| Product Note | This antibody stains the extracellular domain of CD73. The monoclonal antibody 4G4 recognizes both membrane bound and soluble human CD73, also known as ecto-5'-nucleotidase. The monoclonal antibody 4G4 causes a reduction in CD73 expression on lymphocytes, reduces enzyme activity, and inhibits the binding of lymphocytes to endothelial cells. |

| Properties | |
|---------------|---|
| Form | Liquid |
| Buffer | Filter-sterilized PBS, 0.1% BSA |
| Preservative | No preservatives |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. |
| Concentration | $100 \mu g/ml$ (Please refer to the vial label for the specific concentration.) |



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 13 Page 1 of 2



| Immunogen | Inflamed synovial stroma from rheumatoid arthritis patients |
|--------------|---|
| Purification | Protein G purified |
| Endotoxin | < 24 EU/mg (Determined by LAL assay) |
| Conjugation | Unconjugated |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 13 Page 2 of 2