# Src (phospho Tyr416) antibody (Alexa Fluor 488)

## Cat. No. GTX54529

Host	Rabbit	<mark>Package</mark> 100 μl
Clonality	Polyclonal	·
lsotype	lgG	
Applications	WB, ICC/IF	
Reactivity	Human, Mouse, Dog, Chicken	

### Applications

#### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
WB	Assay dependent	
ICC/IF	Assay dependent	
Not tested in other applie	cations.	
Calculated MW	60 kDa. ( <u>Note</u> )	
Product Note	Based on sequence homology, this antibody may cross with other Src family members such as YES1 , FYN proteins when phosphorylated at the equivalent sites.	
Properties		
Form	Liquid	
Buffer	PBS, 0.02% BSA	
Preservative	0.05% Sodium azide	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.	
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)	
Immunogen	This antibody was produced against a chemically synthesized phosphopeptide derived from the region of Src that contains tyrosine 416. The antibody was affinity purified by sequential epitope-specific chromatography, then conjugated to Alexa Fluor 488. The Tyr418 of human Src is equivalent to Tyr416 of chicken Src.	
Purification	Purified IgG	
Conjugation	Alexa Fluor 488 <u>Wavelength</u>	



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 02 Page 1 of 2




Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

#### DATA IMAGES



#### GTX54529 ICC/IF Image

ICC/IF analysis of MDCK cells were left untreated (A), stimulated with H2O2 (1 mM for 10 minutes) (B), or pretreated with PP2 prior to H2O2 stimulation (C) using GTX54529 Src (phospho Tyr416) antibody (Alexa Fluor 488).



For full product information, images and publications, please visit our <u>website</u>.