

TRPC6 antibody

Cat. No. GTX54757

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-Fr, FCM, IP
Reactivity	Human, Mouse, Rat

Package

50 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-Fr	Assay dependent
FCM	Assay dependent
IP	Assay dependent

Not tested in other applications.

Calculated MW 107 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide (C)RRNESQDYLLMDELG, corresponding to amino acid residues 24-38 (Intracellular, N-terminus) of mouse TRPC6 (Accession Q61143).
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX54757 IHC-Fr Image

IHC-Fr analysis of mouse cerebellum tissue using GTX54757 TRPC6 antibody.

Panel A : TRPC6 (red) appears in Purkinje cells (arrows) and in the molecular (Mol) layer.

Panel B : Staining with mouse anti parvalbumin in the same brain section.

Panel C : Merge of TRPC6 and PV demonstrates partial co-localization in the Purkinje and the molecular layers.



GTX54757 WB Image

WB analysis of rat brain membrane lysate using GTX54757 TRPC6 antibody preincubated with or without immunogen peptide.

Dilution : 1:200



For full product information, images and publications, please visit our [website](#).