

MED1 antibody

Cat. No. GTX55164

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-P
Reactivity	Human

Package 100 μΙ

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500 - 1:1000
ICC/IF	1:100 - 1:500
IHC-P	1:100 - 1:200
Not tested in other applications	

Calculated MW 168 kDa. (<u>Note</u>)


PROPERTIES	
Form	Liquid
Buffer	0.42% Potassium Phosphate, 0.87% NaCl, 30% Glycerol
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of MED1. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.


Date 2024 / 05 / 02 Page 1 of 2

DATA IMAGES




GTX55164 WB Image

WB analysis of HeLa (A), HUVEC (B), Jurkat (C) whole cell lysates using GTX55164 MED1 antibody.



GTX55164 IHC-P Image

IHC-P analysis of formalin fixed human breast cancer tissue section using GTX55164 MED1 antibody. Antigen retrieval: Heat mediated antigen retrieval with sodium citrate buffer (pH 6.0)



GTX55164 ICC/IF Image

ICC/IF analysis of formalin-fixed HUVEC cells using GTX55164 MED1 antibody.

Red: Primary antibody

Blue: DAPI

Permeabilization: 0.1% Triton X-100 in TBS for 5-10 minutes



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 02 Page 2 of 2