

Coxsackie Adenovirus Receptor antibody

Cat. No. GTX56276

| Host | Rabbit |
|--------------|-------------------|
| Clonality | Polyclonal |
| Isotype | lgG |
| Applications | WB, ICC/IF |
| Reactivity | Human, Mouse, Rat |

Package 100 μΙ

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500 - 1:1000 |
| ICC/IF | 1:100 - 1:500 |
| | |

Not tested in other applications.

Calculated MW 40 kDa. (Note)

| Properties | |
|---------------|--|
| Form | Liquid |
| Buffer | 0.42% Potassium Phosphate, 0.87% NaCl, 30% Glycerol |
| Preservative | 0.01% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | Batch dependent (Please refer to the vial label for the specific concentration.) |
| Immunogen | KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of Coxsackie Adenovirus Receptor. The exact sequence is proprietary. |
| Purification | Purified by antigen-affinity chromatography |
| Conjugation | Unconjugated |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |




For full product information, images and publications, please visit our website.

Date 2025 / 12 / 28 Page 1 of 2




DATA IMAGES



GTX56276 WB Image

WB analysis of HeLa (A), NIH3T3 (B), H9C2 (C) whole cell lysates using GTX56276 Coxsackie Adenovirus Receptor antibody.



GTX56276 ICC/IF Image

ICC/IF analysis of formalin-fixed NIH3T3 cells using GTX56276 Coxsackie Adenovirus Receptor antibody.

Red: Primary antibody

Blue : DAPI

Permeabilization: 0.1% Triton X-100 in TBS for 5-10 minutes



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 28 Page 2 of 2