

# Histone H4K20me1 (monomethyl Lys20) antibody

# Cat. No. GTX57170

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-P, Dot, ChIP assay
Reactivity	Human, Mouse, Rat

Package 100 μl

# APPLICATION

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500 - 1:2000
ICC/IF	1:50 - 1:200
IHC-P	1:50 - 1:200
Dot	Assay dependent
ChIP assay	1:20 - 1:100

Not tested in other applications.

PROPERTIES	
Form	Liquid
Buffer	PBS, 50% Glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide of human MonoMethyl-Histone H4-K20
Purification	Purified by affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 18 Page 1 of 2




## DATA IMAGES




## GTX57170 ICC/IF Image

ICC/IF analysis of 293T cells using GTX57170 Histone H4K20me1 (monomethyl Lys20) antibody.




## GTX57170 Dot Image

Dot blot analysis of all sorts of methylation peptides using GTX57170 Histone H4K20me1 (monomethyl Lys20) antibody.



#### GTX57170 IHC-P Image

IHC-P analysis of rat testis tissue using GTX57170 Histone H4K20me1 (monomethyl Lys20) antibody. Dilution: 1:200



## GTX57170 WB Image

WB analysis of various sample lysates using GTX57170 Histone H4K20me1 (monomethyl Lys20) antibody.

Dilution : 1:1000 Loading : 25µg per lane



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 18 Page 2 of 2