HDAC3 antibody [GT9080]

Cat. No. GTX60832

| Host | Mouse |
|--------------|--------------------|
| Clonality | Monoclonal |
| lsotype | IgG |
| Applications | ICC/IF, ChIP assay |
| Reactivity | Human |
| | |

Package 50 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| ICC/IF | Assay dependent |
| ChIP assay | Assay dependent |

Not tested in other applications.


| Properties | |
|--------------|--|
| Form | Liquid |
| Buffer | Tissue culture supernatant |
| Preservative | 0.05% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Immunogen | Human HDAC3 (Histone deacetylase 3), using a KLH-conjugated synthetic peptide containing a sequence from the C- terminal region of the protein. |
| Purification | Unpurified |
| Conjugation | Unconjugated |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our <u>website</u>.


Date 2025 / 06 / 30 Page 1 of 2

DATA IMAGES



GTX60832 ICC/IF Image

ICC/IF analysis of HeLa cells using HDAC3 antibody (red) and DAPI. Cells were fixed with 4% formaldehyde for 10' and blocked with PBS/TX-100 containing 5% normal goat serum and 1% BSA. The cells were immunofluorescently labelled with the HDAC3 antibody at a dilution of 1:500.



GTX60832 ChIP assay Image

ChIP was performed with HeLa extracts and either 5 μ g of HDAC3 antibody or control IgG. The precipitated DNA was detected by qPCR with primers targeting to GAPDH promoter, c-fos promoter and the coding region of p21, a known target gene of HDAC3.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 30 Page 2 of 2