

## GLP1 antibody [10] (Biotin)

**Cat. No. GTX60952-02**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2a
<b>Applications</b>	ELISA
<b>Reactivity</b>	Human

**Package**  
50 µl

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ELISA	Assay dependent

Not tested in other applications.

**Product Note**

This antibody binds the free N-terminus of GLP-1(7-37) and GLP-1(7-36)amide and shows <0.2% cross-reactivity with GLP-1(9-36)amide, glucagon and human GIP. Cross-reacts 0.25% with GLP-1(1-37), approximately 6% with human GLP-2 and approximately 50% with exendin-4. this antibody cross-reacts with Liraglutide.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	10mM Phosphate, 140mM NaCl
<b>Preservative</b>	15mM Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Synthetic GLP-1(7-17)
<b>Purification</b>	Biotinylated
<b>Conjugation</b>	Biotin

**Note**


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



**GTX60952-02 ELISA Image**

The calibration curve of a sandwich assay for Non-amidated GLP-1 using GTX60954 as the capture antibody and GTX60952 -02 as the biotinylated detection antibody.



**GTX60952-02 ELISA Image**

This uses GTX60951 as capture antibody in combination with biotinylated detection antibody GTX60952 -02 exclusively to measure GLP-1(7-36)amide in biological samples. GLP-1(7-36)amide is the principal active form of the peptide, and has been estimated to make up about 70% of the physiological response. Relative to measuring both active forms of GLP-1 together, the responses are reduced by not measuring the approximately 30% contribution of GLP-1(7-37) to the overall response. There is no cross-reactivity with any extended or truncated physiological form of GLP-1(7-36)amide. Specimens must be collected into tubes containing an adequate amount of DPP-4 inhibitor (in addition to routine peptidase inhibitors) to prevent the rapid in vitro degradation of the active peptide.



For full product information, images and publications, please visit our [website](#).