

HMGA2 antibody [GT763]

Cat. No. GTX629478

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF, IHC-P, IP
Reactivity	Human, Mouse, Rat

References (6)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500

Not tested in other applications.


Calculated MW 12 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human HMGA2. The exact sequence is proprietary.
Purification	Affinity purified by Protein G.
Conjugation	Unconjugated
Note	For <i>In vitro</i> laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Due to intellectual property restrictions, this product is not available for purchase in AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE.



For full product information, images and publications, please visit our [website](#).


DATA IMAGES

GTX629478 IHC-P Image

HMGA2 antibody [GT763] detects HMGA2 protein at nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded mouse brain.

HMGA2 stained by HMGA2 antibody [GT763] (GTX629478) diluted at 1:200.


Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTX629478 IP Image

Immunoprecipitation of HMGA2 protein from HepG2 nuclear extracts using 5 µg of HMGA2 antibody [GT763] (GTX629478).

Western blot analysis was performed using HMGA2 antibody [GT763] (GTX629478) diluted at 1:500.

EasyBlot anti-Rabbit IgG (GTX221666-01) was used as a secondary reagent.



GTX629478 IHC-P Image

HMGA2 antibody [GT763] detects HMGA2 protein on C2C12 by immunohistochemical analysis.

Sample: Paraffin-embedded C2C12.

HMGA2 antibody [GT763] (GTX629478) dilution: 1:200.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min


GTX629478 WB Image

Whole cell extract (30 µg) was separated by 15% SDS-PAGE, and the membrane was blotted with HMGA2 antibody [GT763] (GTX629478) diluted at 1:1000. The HRP-conjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).