

5-Hydroxymethylcytosine / 5-hmC antibody [GT13612]

Cat. No. GTX629765

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Application	IHC-P, IHC-Fr, Dot, EM, MeDIP
Reactivity	Species independent

Reference (9)

★★★★★ Review (1)

Package

100 µl, 25 µl

PRODUCT

Summary

5-hmC antibody recognizes 5-Hydroxymethylcytosine (5-hmC), which is referred to as a "sixth base" of mammalian genomic DNA. 5-hmC is thought to be an epigenetic regulatory modification in its own right, while also being the product of the first TET protein-catalyzed oxidation of 5-methylcytosine in the DNA demethylation pathway.

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100-1:1000
IHC-Fr	Assay dependent
Dot	Assay dependent
EM	Assay dependent
MeDIP	Assay dependent

Not tested in other applications.

PROPERTIES

Form	Liquid
Buffer	PBS
Preservative	No preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.81 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	The immunogen used to generate this antibody corresponds to 5-HydroxymethylCytosine / 5-hmC
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Date 2024 / 05 / 04 Page 1 of 2


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

GTX629765 MeDIP Image

GTX629765 5-HydroxymethylCytidine antibody [GT13612] in MeDIP experiment.
Human genomic DNA (500 ng) mixed with methylated cytosine standard kit (GTX400004) were subjected to MeDIP with 5-HydroxymethylCytidine antibody [GT13612] (GTX629765) at dilution of 1:500 and analyzed with semi-quantitative PCR. The immunoprecipitated DNA was plotted as % of input DNA.



GTX629765 IHC-P Image

5-HydroxymethylCytidine antibody [GT13612] detects 5-HydroxymethylCytidine protein at nucleus on mouse testis by immunohistochemical analysis.

Sample: Paraffin-embedded mouse testis.

5-HydroxymethylCytidine antibody [GT13612] (GTX629765) dilution: 1:100.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX629765 IHC-P Image

5-HydroxymethylCytidine antibody [GT13612] detects 5-HydroxymethylCytidine protein at nucleus on mouse prostate by immunohistochemical analysis.

Sample: Paraffin-embedded mouse prostate.

5-HydroxymethylCytidine antibody [GT13612] (GTX629765) dilution: 1:100.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min


GTX629765 Dot Image

GTX629765 5-hydroxymethylcytosine antibody [GT13612] Dot blot analysis of anti-5-hmC antibody with the synthetic DNA controls (GTX400004). DNA samples (0.5 to 50 ng as indicated) were spotted onto the positively charged Nylon membrane and blotted with 5-hmC antibody (GTX629765) at a dilution of 1:500.
A: Unmethylated DNA fragment B: DNA fragment containing 5-methylcytosine C: DNA fragment containing 5-hydroxymethylcytosine D: DNA fragment containing 5-formylcytosine E: DNA fragment containing 5-carboxylcytosine



For full product information, images and publications, please visit our [website](https://www.genetex.com).