SQSTM1 / P62 antibody [GT1478]

Cat. No. GTX629890

Host	Mouse	References (10)
Clonality	Monoclonal	🚖 🚖 🚖 🌟 🏌 Review (9)
lsotype	lgG2b	<mark>Package</mark> 100 μl, 25 μl
Applications	WB, ICC/IF, IHC-Fr, FCM	
Reactivity	Human, Mouse	

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
WB	1:500-1:3000	
ICC/IF	1:100-1:1000	
IHC-Fr	1:100-1:1000	
FCM	1:50-1:200	
Not tested in other applications		

Not tested in other applications.

Calculated MW

48 kDa. (<u>Note</u>)

Properties		
Form	Liquid	
Buffer	PBS, 20% Glycerol	
Preservative	No preservatives	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen	Recombinant protein encompassing a sequence within the center region of human SQSTM1 / P62. The exact sequence is proprietary.	
Purification	Affinity purified by Protein G.	
Conjugation	Unconjugated	



For full product information, images and publications, please visit our website.


Date 2025 / 09 / 04 Page 1 of 2

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX629890 WB Image

SQSTM1 antibody detects SQSTM1 protein by western blot analysis. Un-treated (-) and treated (+, Thapsigargin treatment for 12 hr and 24hr) Huh-7 whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 antibody (GTX629890) diluted by 1:1000. The ACTB was used as internal control (GTX110564, 1:50000) shown at the bottom panel. The HRP-conjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody.



GTX629890 WB Image

Various whole cell extracts (30 μ g) were separated by 10% SDS-PAGE, and the membrane was blotted with SQSTM1 / P62 antibody [GT1478] (GTX629890) diluted at 1:500. The HRP-conjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody.



GTX629890 FCM Image

SQSTM1 antibody [GT1478] (GTX629890) detects SQSTM1 protein by flow cytometry analysis. Sample: HeLa cell fixed in 4% paraformaldehyde at 4oC for 5 min. Brown: Unlabelled sample was also used as a control. Blue: SQSTM1 antibody [GT1478] (GTX629890) dilution: 1:100. Acquisition of >20,000 events were collected using Argon ion laser (488nm) and 525/30 bandpass filter.



For full product information, images and publications, please visit our <u>website</u>.