

# NDP52 antibody [GT1813]

# Cat. No. GTX630397

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2a
Applications	WB, ICC/IF
Reactivity	Human

Package 100 μl, 25 μl

# Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000

Not tested in other applications.

Calculated MW 52 kDa. (Note)


Properties	
Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human NDP52. The exact sequence is proprietary.
Purification	Affinity purified by Protein G.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.


Date 2025 / 11 / 03 Page 1 of 2

## DATA IMAGES



## GTX630397 WB Image


NDP52 antibody detects NDP52 protein by western blot analysis. Un-treated (-) and treated (+, Thapsigargin treatment for 12hrs and 24hrs) Huh-7 whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with NDP52 antibody (GTX630397) diluted by 1:500. The ACTB was used as internal control (GTX110564, 1:50000) shown at the bottom panel.



#### GTX630397 ICC/IF Image


NDP52 antibody [GT1813] detects NDP52 protein at autophagosome by immunofluorescent analysis. Samples: HeLa cells mock (left) and treated with  $50\mu M$  Chloroquine for 24 hr (right) were fixed in 4% paraformaldehyde at RT for 15 min.

Green: NDP52 protein stained by NDP52 antibody [GT1813] (GTX630397) diluted at 1:1000. Red: Phalloidin, a F-actin marker.



### GTX630397 WB Image

Non-transfected (–) and transfected (+) HepG2 whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with NDP52 antibody [GT1813] (GTX630397) diluted at 1:1000.



#### GTX630397 WB Image

NDP52 antibody detects NDP52 protein by western blot analysis. Un-treated (-) and treated (+, Thapsigargin treatment for 12hrs and 24hrs) HepG2 whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with NDP52 antibody (GTX630397) diluted by 1:500. The ACTB was used as internal control (GTX110564, 1:50000) shown at the bottom panel.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 03 Page 2 of 2