Histone H4K20me2 (dimethyl Lys20) antibody [GT1851]

Cat. No. GTX630545

Clonality Mor	noclonal
lsotype lgG	1
Application WB,	, ICC/IF, IP, Dot, ChIP assay
Reactivity Hun	nan, Mouse

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:10000
ICC/IF	1:100-1:1000
IP	1:100-1:500
Dot	1:500
ChIP assay	Assay dependent

Reference (4) Package 100 μl, 25 μl

Not tested in other applications.

PROPERTIES	
Form	Liquid
Buffer	PBS
Preservative	No preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide surrounding di-methyl Lys20 of human Histone H4. The exact sequence is proprietary.
Purification	Affinity purified by Protein G.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.
-	




For full product information, images and publications, please visit our <u>website</u>.

DATA IMAGES

(ng)	Un	Me	2Me	3Me	
50	0				H4 antibody (GTX129560)
50		0			H4K20me1 antibody (GTX128959)
50 0					H4K20me2 antibody (GTX630545)
50				0	H4K20me3 antibody (GTX128960)


GTX630545 Dot Image

Dot blot analysis of various of H4K20 methylation peptides using Histone H4K20me2 (dimethyl Lys20) antibody [GT1851] (GTX630545) at antibody dilution of 1:500. Un : H4 protein peptide Me : H4K20 mono-methyl peptide 2Me : H4K20 di-methyl peptide 3Me : H4K20 Tri-methyl peptide



GTX630545 ChIP assay Image

Cross-linked ChIP was performed with HeLa chromatin extract and 5 μ g of either control mouse IgG or anti-Histone H4K20me2 (dimethyl Lys20) antibody. The precipitated DNA was detected by PCR with primer set targeting to MyoD.



GTX630545 WB Image


Histone H4 (di methyl Lys20) antibody [GT1851] detects Histone H4 (di methyl Lys20) protein by western blot analysis.

A. 30 µg HeLa whole cell lysate/extract

B. 30 µg HeLa nuclear lysate/extract

Histone H4 (di methyl Lys20) antibody [GT1851] (GTX630545) dilution: 1:5000

The HRP-conjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody.



GTX630545 IP Image

Immunoprecipitation of HIST1H4A protein from HeLa whole cell extracts using 5 μ g of Histone H4K20me2 (dimethyl Lys20) antibody [GT1851] (GTX630545).

Western blot analysis was performed using Histone H4K20me2 (dimethyl Lys20) antibody [GT1851] (GTX630545).

EasyBlot anti-mouse IgG (GTX221667-01) was used as a secondary reagent.



For full product information, images and publications, please visit our <u>website</u>.