

BICD2 antibody [GT1824]

Cat. No. GTX631995

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Application	WB, IP
Reactivity	Human, Rat

Package 100 μl, 25 μl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IP	1:100-1:500

Not tested in other applications.

Calculated MW 94 kDa. (Note)

PROPERTIES	
Form	Liquid
Buffer	PBS
Preservative	No preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human BICD2. The exact sequence is proprietary.
Purification	Affinity purified by Protein G.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 05 Page 1 of 2

€ 886-3-6208988 📻 886-3-6208989 🐷 infoasia@genetex.com




DATA IMAGES




GTX631995 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with BICD2 antibody [GT1824] (GTX631995) diluted at 1:500.



GTX631995 WB Image

BICD2 antibody detects BICD2 protein by western blot analysis. Various whole cell extracts (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with BICD2 antibody (GTX631995) diluted by 1:1000.



GTX631995 WB Image

Various whole cell extracts (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with BICD2 antibody [GT1824] (GTX631995) diluted at 1:500.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 05 Page 2 of 2