

DCK antibody [GT10412]

Cat. No. GTX632065

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2b
Applications	WB, ICC/IF, IHC-P, IP
Reactivity	Human, Mouse, Rat

References (1) Package 100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:10000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500

Not tested in other applications.

Calculated MW 31 kDa. (Note)


Properties	
Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	2.01 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human DCK. The exact sequence is proprietary.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.


Date 2026 / 01 / 02 Page 1 of 2

DATA IMAGES




GTX632065 WB Image

DCK antibody [GT10412] detects DCK protein by western blot analysis. Various whole cell extracts (30 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody [GT10412] (GTX632065) diluted at a dilution of 1:500.



GTX632065 WB Image

DCK antibody detects DCK protein by western blot analysis. Various whole cell extracts (30 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody (GTX632065) diluted at a dilution of 1:500.



GTX632065 WB Image

DCK antibody [GT10412] detects DCK protein by western blot analysis. Various whole cell extracts (30 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody [GT10412] (GTX632065) diluted at a dilution of 1:500.



For full product information, images and publications, please visit our <u>website</u>.

Date 2026 / 01 / 02 Page 2 of 2