

Venezuelan Equine Encephalitis Virus nsP2 antibody [HL1919]

Cat. No. GTX637668

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF
Reactivity	Venezuelan equine encephalitis virus

References (1)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	1:100-1:1000

Not tested in other applications.

Product Note

This antibody was raised against Venezuelan Equine Encephalitis Virus nsP2, and it does not cross-react with Western Equine Encephalitis Virus and Eastern Equine Encephalitis Virus.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of Venezuelan Equine Encephalitis Virus nsP2. (Strain P676)
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX637668 ICC/IF Image

Immunofluorescent analysis of Venezuelan equine encephalitis virus infected cells using Venezuelan Equine Encephalitis Virus nsP2 antibody [HL1919] (GTX637668).

Sample: Multiple virus infected cells slide.

Green: Venezuelan Equine Encephalitis Virus nsP2 antibody [HL1919] (GTX637668) diluted at 1:100.


Blue: Fluoroshield with DAPI (GTX30920).



GTX637668 WB Image

Non-transfected (-) and transfected (+) 293T whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Venezuelan Equine Encephalitis Virus nsP2 antibody [HL1919]

(GTX637668) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX637668 WB Image

Non-transfected (-) and transfected (+) 293T whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Venezuelan Equine Encephalitis Virus nsP2 antibody [HL1919]

(GTX637668) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](http://www.genetex.com).