

HHIP antibody [HL1969]

Cat. No. GTX637862

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	Assay dependent

Not tested in other applications.

Observed MW (kDa) 79 kDa.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of human HHIP
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


DATA IMAGES

GTx637862 IHC-P Image

HHIP antibody [HL1969] detects HHIP protein at cell membrane and secreted by immunohistochemical analysis.

Sample: Paraffin-embedded mouse kidney.

HHIP stained by HHIP antibody [HL1969] (GTx637862) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTx637862 IHC-P Image

HHIP antibody [HL1969] detects HHIP protein at cell membrane and secreted by immunohistochemical analysis.

Sample: Paraffin-embedded mouse duodenum.

HHIP stained by HHIP antibody [HL1969] (GTx637862) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTx637862 IHC-P Image

HHIP antibody [HL1969] detects HHIP protein at cell membrane and secreted by immunohistochemical analysis.


Sample: Paraffin-embedded human hepatocellular carcinoma.

HHIP stained by HHIP antibody [HL1969] (GTx637862) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTx637862 WB Image

Various whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with HHIP antibody [HL1969] (GTx637862) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced. Corresponding RNA expression data for the same cell lines are based on Human Protein Atlas program.


GTx637862 WB Image

Non-transfected (-) and transfected (+) 293T whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with HHIP antibody [HL1969] (GTx637862) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).