

ULK2 antibody [HL2206]

Cat. No. GTX638210

Host	Rabbit
Clonality	Monoclonal
Isotype	lgG
Applications	WB, ICC/IF
Reactivity	Human, Rat

Package 100 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
Niet tested in ethan and it estima	

Not tested in other applications.

Observed MW (kDa) 113 kDa.

Properties	
Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragmemt of human ULK2
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 28 Page 1 of 2




DATA IMAGES




GTX638210 WB Image

Whole cell extract (30 µg) was separated by 7.5% SDS-PAGE, and the membrane was blotted with ULK2 antibody [HL2206] (GTX638210) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX638210 ICC/IF Image

ULK2 antibody [HL2206] detects ULK2 protein at cytoplasm and nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: ULK2 stained by ULK2 antibody [HL2206] (GTX638210) diluted at 1:500. Blue: Fluoroshield with DAPI (GTX30920).



GTX638210 WB Image

Various whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with ULK2 antibody [HL2206] (GTX638210) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 28 Page 2 of 2