

COX4 antibody [HL2264]

Cat. No. GTX638316

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat, Zebrafish

Package 100 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	Assay dependent
Not tested in other applications.	

Observed MW (kDa) 17 kDa.

Properties	
Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of human COX4.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 27 Page 1 of 2




DATA IMAGES




GTX638316 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 15% SDS-PAGE, and the membrane was blotted with COX4 antibody [HL2264] (GTX638316) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX638316 WB Image

Various whole cell extracts (30 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with COX4 antibody [HL2264] (GTX638316) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX638316 IHC-P Image

COX4 antibody [HL2264] detects COX4 protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse tissues.

COX4 stained by COX4 antibody [HL2264] (GTX638316) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX638316 ICC/IF Image

COX4 antibody [HL2264] detects COX4 protein at mitochondria by immunofluorescent analysis. Sample: HeLa cells were fixed in ice-cold MeOH for 5 min.

Green: COX4 stained by COX4 antibody [HL2264] (GTX638316) diluted at 1:500.

Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:1000.

Blue: Fluoroshield with DAPI (GTX30920).



GTX638316 WB Image

Various tissue extracts (50 μ g) were separated by 15% SDS-PAGE, and the membrane was blotted with COX4 antibody [HL2264] (GTX638316) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 27 Page 2 of 2