

NKG2A antibody [HL2718]

Cat. No. GTX639545

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	Assay dependent

Not tested in other applications.

Observed MW (kDa) 38-40 kDa.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide of human NKG2A.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX639545 WB Image

Whole cell extract (30 µg) was separated by 12% SDS-PAGE, and the membrane was blotted with NKG2A antibody [HL2718] (GTX639545) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX639545 IHC-P Image

NKG2A antibody [HL2718] detects NKG2A protein by immunohistochemical analysis.

Sample: Paraffin-embedded human spleen.

NKG2A stained by NKG2A antibody [HL2718] (GTX639545) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX639545 ICC/IF Image

NKG2A antibody [HL2718] detects NKG2A protein by immunofluorescent analysis.

Sample: NK-92 cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: NKG2A stained by NKG2A antibody [HL2718] (GTX639545) diluted at 1:100.

Blue: Fluoroshield with DAPI (GTX30920).



For full product information, images and publications, please visit our [website](https://www.genetex.com).