

SCAP antibody [HL2796]

Cat. No. GTX639684

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse, Rat

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	Assay dependent

Not tested in other applications.

Observed MW (kDa) 120, 125 kDa.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of human SCAP.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX639684 WB Image

Various whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with SCAP antibody [HL2796] (GTX639684) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX639684 IHC-P Image

SCAP antibody [HL2796] detects SCAP protein at cytoplasm by immunohistochemical analysis. Sample: Paraffin-embedded rat stomach.

SCAP stained by SCAP antibody [HL2796] (GTX639684) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX639684 IHC-P Image

SCAP antibody [HL2796] detects SCAP protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded mouse colon.

SCAP stained by SCAP antibody [HL2796] (GTX639684) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX639684 WB Image

Various whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with SCAP antibody [HL2796] (GTX639684) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).