

AMFR antibody [HL2865]

Cat. No. GTX640142

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse, Rat

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	Assay dependent

Not tested in other applications.

Observed MW (kDa) 73 kDa.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment of Human AMFR.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX640142 WB Image

Boiled and unboiled mouse tissue extract (50 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with AMFR antibody [HL2865] (GTX640142) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX640142 WB Image

Boiled and unboiled rat tissue extract (50 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with AMFR antibody [HL2865] (GTX640142) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX640142 WB Image

Boiled and unboiled HepG2 whole cell and membrane extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with AMFR antibody [HL2865] (GTX640142) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody. (WCE: whole cell extract; ME: membrane extract)



GTX640142 IHC-P Image

AMFR antibody [HL2865] detects AMFR protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse liver.

AMFR stained by AMFR antibody [HL2865] (GTX640142) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 20 min



For full product information, images and publications, please visit our [website](https://www.genetex.com).