

DDDDK tag antibody [M2-RB]

Cat. No. GTX640415

Host	Rabbit
Clonality	Monoclonal
Isotype	lgG
Applications	WB, ICC/IF, ELISA
Reactivity	Species independent

Package $100~\mu\text{l},\,25~\mu\text{l}$

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:10000
ICC/IF	Assay dependent
ELISA	Assay dependent

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	The immunogen used to generate this antibody corresponds to DDDDK tag.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 11 / 03 Page 1 of 2




DATA IMAGES



GTX640415 WB Image


Non-transfected (–) and transfected (+) 293T whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membranes were blotted with DDDDK tag antibody [M2-RB] (GTX640415) diluted at 1:10000 and competitor's antibody (Highly Cited Antibody) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

*The competitor is not affiliated with GeneTex and does not endorse this product.




GTX640415 ELISA Image

Indirect ELISA analysis was performed by coating the plate with recombinant Multiple Tags Positive Control (6xHis-MBP-DDDDK-V5-HSV-HA-T7-Myc-S) (GTX130342-pro) (57.14-0.89 nM). Coated protein was probed with DDDDK tag antibody [HL3005] (GTX640415) ($1 \mu g/mL$). Goat anti-rabbit IgG antibody (HRP) (GTX213110-01) (1:10000) was used to detect the bound primary antibody.




GTX640415 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 10% SDS-PAGE, and the membrane was blotted with DDDDK tag antibody [M2-RB] (GTX640415) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX640415 ELISA Image

Indirect ELISA analysis was performed by coating the plate with recombinant Multiple Tags Positive Control (GST-DDDDK-V5-HSV-HA-T7-Myc-S) (GTX130343-pro) (72.73-1.14 nM). Coated protein was probed with DDDDK tag antibody [HL3005] (GTX640415) (1 μ g/mL). Goat anti-rabbit IgG antibody (HRP) (GTX213110-01) (1:10000) was used to detect the bound primary antibody.



GTX640415 ICC/IF Image

DDDDK tag antibody [M2-RB] detects DDDDK tag protein by immunofluorescent analysis. Sample: Mock and transfected 293T cells were fixed in ice-cold MeOH for 5 min. Green: DDDDK tag stained by DDDDK tag antibody [M2-RB] (GTX640415) diluted at 1:500. Blue: Fluoroshield with DAPI (GTX30920).



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 03 Page 2 of 2