Hepatitis C virus Core Antigen antibody [H6-29] (FITC)

Cat. No. GTX64115

Host	Mouse	<mark>Package</mark> 50 μg
Clonality	Monoclonal	
lsotype	lgG2a	
Application	WB, ICC/IF, IHC	
Reactivity	Hepatitis C virus	

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC	Assay dependent


Not tested in other applications.

PROPERTIES		
Form	Liquid	
Buffer	Filter-sterilized PBS, 50% Glycerol	
Preservative	No preservative	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.	
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)	
Immunogen	A part of the core region (nucleotides 369-704, amino acids 13-124) of HCV genotype 1b expressed in E. coli	
Conjugation	Fluorescein isothiocyanate (FITC)	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
NOLE	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	




For full product information, images and publications, please visit our <u>website</u>.

DATA IMAGES




GTX64115 IHC Image

Immunohistochemical detection of HCV core protein. Tissue section from a patient with chronic hepatitis C was immunostained to reveal cells expressing HCV core antigen, which are scattered in the lobules (indirect immunohistochemical method, counterstained with Mayer's hematoxylin).



GTX64115 WB Image

Western blotting of HCV core protein. Chimp liver cells were infected with recombinant vaccinia virus containing a HCV genome cDNA and were subjected to Western blotting using this antibody. The core protein is detected as a 22-kDa band.



GTX64115 ICC/IF Image

Detection of HCV core protein by immunofluorescence antibody staining. Chimp liver cells were infected with recombinant vaccinia virus containing a HCV genome cDNA. After incubation for 48 hr, the cells were fixed with acetone and HCV core protein was detected by indirect immunofluorescence staining using this antibody.



For full product information, images and publications, please visit our <u>website</u>.