

# Mouse GITRL protein, DDDDK tag (active)

**Cat. No. GTX65629-pro**

**Applications** Functional Assay

**Species** Mouse

**Package**

50 µg

## Applications

### Application Note

Induces GTR-mediated NF-κB activation in HEK 293 cells.

**Product Note** Binds to mouse GTR.

## Properties

**Form** Liquid

**Buffer** 10mM Glycine

**Preservative** No preservatives

**Storage** Store as concentrated solution. Aliquot and store at -20°C or below. Avoid freeze-thaw cycles.

**Concentration** 0.5 mg/ml (Please refer to the vial label for the specific concentration.)

**Region/Sequence** N-terminal DDDDK-Tag; the extracellular domain of mouse GITRL (a.a. 45-173).

**Expression System** HEK293 cells


**Purity** ≥90% by SDS-PAGE.

**Endotoxin** < 0.1 EU/µg

**Conjugation** Unconjugated

**Note** For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

## DATA IMAGES



### GTX65629-pro Image

GTX65629-pro GITRL protein induced GTR-mediated NF-κB activation. HEK293 cells were transiently transfected with 20ng of a (NF-κB)-Luc reporter, 2ng of a Renilla luciferase, and with 20ng of a GITR full length encoding construct. At 24hr after transfection, the indicated amounts of GITRL protein or a DDDDK control protein were treated for 4hr and followed by the measurement of dual luciferase activities.



For full product information, images and publications, please visit our [website](#).