

Human IL2 (Superkine) protein, human IgG2 Fc tag (active)

Cat. No. GTX65631-pro**Applications** Functional Assay**Species** Human**Package**

10 µg

Applications

Application Note


Functional assay: Triggers far greater antitumor responses than native IL-2 in vivo but with lower toxicity.

Product Note Binds to human and mouse IL-2R.

Properties

Form Lyophilized powder**Buffer** Reconstitute with 100µl ddH₂O to 0.1mg/ml. Lyophilized from 0.2µm-filtered PBS.**Preservative** No preservatives**Storage** Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.**Region/Sequence** The extracellular domain of human IL-2 superkine (a.a. 21-153) (mutant H9 containing the mutations L80F / R81D / L85V / I86V / I92F) is fused at the C-terminus to the Fc portion of human IgG2.**Expression System** HEK293 cells**Purity** ≥95% by SDS-PAGE.**Endotoxin** < 0.01 EU/µg**Conjugation** Unconjugated**Note** For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

DATA IMAGES

**GTX65631-pro Image**

Binding of GTX65631-pro IL2 (Superkine) human Fc fusion protein to IL-2R β (human) is increased > 10 fold compared to IL-2 human Fc fusion protein. Method: IL-2R β (human) protein was coated on an ELISA plate at 1µg/ml. After blocking and washing steps, indicated concentrations of GTX65631-pro IL-2 Superkine (Fc) or IL-2 (Fc) protein were added. Following incubation for 1 h at RT, the binding was detected using an anti-human Fc antibody (HRP).



For full product information, images and publications, please visit our [website](#).