

# Human CD134 protein, human IgG1 Fc tag (active)

**Cat. No. GTX65665-pro**

<b>Applications</b>	Functional Assay
<b>Species</b>	Human

**Package**  
50 µg

## Applications

### Application Note

Blocks T cell proliferation induced by CD134L.

**Product Note** Binds to human and mouse OX40L.

## Properties

<b>Form</b>	Lyophilized powder
<b>Buffer</b>	Reconstitute with 50µl ddH <sub>2</sub> O to 1mg/ml. Lyophilized from 0.2µm-filtered PBS.
<b>Preservative</b>	No preservatives
<b>Storage</b>	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
<b>Region/Sequence</b>	Human CD134 (a.a. 29-214) is fused at the C-terminus to the Fc portion of human IgG1.
<b>Expression System</b>	HEK293 cells
<b>Purity</b>	??95% by SDS-PAGE.
<b>Endotoxin</b>	< 0.01 EU/µg
<b>Conjugation</b>	Unconjugated
<b>Note</b>	For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).