

Human B7-H4 protein, human IgG1 Fc tag (active)

Cat. No. GTX65697-pro**Applications** Functional Assay**Species** Human**Package**

100 µg

Applications

Application Note

Measured by the ability to inhibit anti-CD3-induced proliferation of stimulated human T cells. Human T lymphocytes cultured for 72 hours with PHA were incubated for an additional 3 days in 96 well plate coated with 500ng/ml anti-CD3 and 10 µg/ml human B7H4/Fc. The presence of human B7H4/Fc at 10 µg/ml inhibited anti CD3 response by 30-50%. Optimal dilutions should be determined by each laboratory for each application.

Properties

Form Lyophilized powder**Buffer** Reconstitute with 100 µl ddH₂O. Lyophilized from 0.2µm-filtered PBS.**Preservative** No preservatives**Storage** Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.**Region/Sequence** The extracellular domain of human B7-H4 (a.a. 29-258) is fused to the N-terminus of the Fc region of human IgG1.**Expression System** CHO cells**Purity** ≥98% by SDS-PAGE.**Endotoxin** < 0.06 EU/µg**Conjugation** Unconjugated**Note** For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

For full product information, images and publications, please visit our [website](#).