

## Human ICOS Ligand protein, mouse IgG2a Fc tag (active)

## Cat. No. GTX65721-pro

<b>Applications</b>	Functional Assay
---------------------	------------------

<b>Species</b>	Human
----------------	-------

## Package

100 µg

## Applications

## Application Note

Measured by its ability to stimulate human T cell proliferation in the presence of anti-CD3.

## Properties

<b>Form</b>	Lyophilized powder
-------------	--------------------

<b>Buffer</b>	Reconstitute with 100 µl ddH <sub>2</sub> O. Lyophilized from 0.2µm-filtered PBS.
---------------	---

<b>Preservative</b>	No preservatives
---------------------	------------------

<b>Storage</b>	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
----------------	---

<b>Region/Sequence</b>	The extracellular domain of human CD275 [B7-H2] (a.a. 1-256) is fused to the N-terminus of the Fc region of mouse IgG2a.
------------------------	--

<b>Expression System</b>	CHO cells
--------------------------	-----------

<b>Purity</b>	≥98% by SDS-PAGE.
---------------	-------------------

<b>Endotoxin</b>	< 0.06 EU/µg
------------------	--------------

<b>Conjugation</b>	Unconjugated
--------------------	--------------

<b>Note</b>	For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
-------------	---



For full product information, images and publications, please visit our [website](#).