

## Human ICOS Ligand protein, mutant human IgG1 Fc tag (active)

## Cat. No. GTX65746-pro

Applications	Functional Assay	Package
Species	Human	100 µg

## PRODUCT

Summary	Non-lytic: Acts as a long lasting fusion protein which only binds to the receptor. Mutations to the complement (C1q) and FcgR I binding sites of the IgGs Fc fragment render the fusion proteins incapable of antibody directed cytotoxicity (ADCC) and complement directed cytotoxicity (CDC).
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

## Applications

## Application Note

Measured by its ability to stimulate human T cell proliferation in the presence of anti-CD3.

## Properties

Form	Lyophilized powder
Buffer	Reconstitute with 100 µl ddH <sub>2</sub> O. Lyophilized from 0.2µm-filtered PBS.
Preservative	No preservatives
Storage	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
Region/Sequence	The extracellular domain of human CD275 [B7-H2] (a.a. 1-256) is fused to the N-terminus of the Fc region of a mutant human IgG1.
Expression System	CHO cells
Purity	≥98% by SDS-PAGE.
Endotoxin	< 0.06 EU/µg
Conjugation	Unconjugated
Note	For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 1 of 1