

## Mouse BTNL2 protein, mouse IgG2a Fc tag (active)

**Cat. No. GTX65764-pro**

<b>Applications</b>	Functional Assay
<b>Species</b>	Mouse

**Package**  
100 µg

## Applications

**Application Note**

A butyrophilin-like molecule that functions to inhibit T cell activation.

## Properties

<b>Form</b>	Lyophilized powder
<b>Buffer</b>	Reconstitute with 0.2µm-filtered PBS to 100 µg/ml. Lyophilized from 0.2µm-filtered PBS.
<b>Preservative</b>	No preservatives
<b>Storage</b>	Store at -20°C or below. After reconstitution, keep as concentrated solution. Aliquot and avoid freeze-thaw cycles.
<b>Region/Sequence</b>	The extracellular domain of mouse BTNL2 (a.a. 26-457) is fused to the N-terminus of the Fc region of mouse IgG2a.
<b>Expression System</b>	CHO cells
<b>Purity</b>	≥98% by SDS-PAGE.
<b>Endotoxin</b>	< 0.06 EU/µg
<b>Conjugation</b>	Unconjugated
<b>Note</b>	For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).