

## Human Cystatin D protein, His tag (active)

Cat. No. GTX66965-pro

**Applications** Functional Assay**Species** Human**Package**

50 µg

## Applications

## Application Note

The IC<sub>50</sub> value is < 2.0 nM. The inhibitory function of Cystatin 5 on protease activity of papain was measured by a fluorometric assay using Z-FR-AMC at pH 7.5 at 25°C.


**Observed MW (kDa)** 13.5-18 kDa.

## Properties

**Form** Liquid**Buffer** PBS, 10% Glycerol**Preservative** No preservative**Storage** Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.**Concentration** 0.5 mg/ml (Please refer to the vial label for the specific concentration.)**Region/Sequence** C-terminal His-Tag; Length: 123 a.a. Sequence: QSRTLGGIHTDLNDKSVQ RALDFAISEY NKVINKDEYY SRPLQVMAAY QQIVGGVNNY FNVKFGRTTC TKSQPNLDNC PFNDQPKLKE EEFCFSQINE VPWEDKISIL NYKCRKVHHH HHH**Expression System** Baculovirus**Purity** > 90% by SDS-PAGE.**Endotoxin** < 1.0 EU/µg (determined by LAL method)**Conjugation** Unconjugated**Note** For In vitro laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

For full product information, images and publications, please visit our [website](#).

## DATA IMAGES



### GTx66965-pro Image

3 µg of GTx66965-pro Human Cystatin D protein (active) by SDS-PAGE under reducing condition and visualized by coomassie blue stain



For full product information, images and publications, please visit our [website](#).