

p53 antibody [DO1]

Cat. No. GTX70214

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2a
Application	WB, ICC/IF, IHC-P, IHC-Fr, FACS, IP, ELISA
Reactivity	Human, Mouse



PRODUCT

Summary

p53 antibody recognizes p53 protein, which has a predicted molecular weight of 44 kDa but is typically detected by western blot at 53 kDa. p53, known widely as a tumor suppressor, plays critical roles in various cellular processes including cell cycle regulation, DNA repair, and apoptosis. p53 is mutated in many human cancers.

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:10000
ICC/IF	Assay dependent
IHC-P	1:1000
IHC-Fr	Assay dependent
FACS	Assay dependent
IP	1:100-1:500
ELISA	Assay dependent

Note: Antigen retrieval can be done with EDTA at pH8.0 - Some customers have stated that the product works well without antigen retrieval.

Not tested in other applications.

Calculated MW 44 kDa. (Note)

PROPERTIES	
Form	Liquid
Buffer	1XPBS (pH7), 20% Glycerol
Preservative	No Preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.25 mg/ml (Please refer to the vial label for the specific concentration.)




For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 27 Page 1 of 2


Immunogen	Recombinant human p53 expressed in E. coli.
Purification	Protein A purified From tissue culture supernatant
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX70214 WB Image

Non-transfected (–) and transfected (+) HCT116 whole cell extracts (30 μ g) were separated by 10% SDS-PAGE, and the membrane was blotted with p53 antibody [DO1] (GTX70214) diluted at 1:1000. The HRP-conjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody.



GTX70214 IHC-P Image

p53 antibody [DO1] detects p53 protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded human pancreatic cancer. p53 stained by p53 antibody [DO1] (GTX70214) diluted at 1:100. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX70214 ICC/IF Image

p53 antibody [DO1] detects p53 protein at nucleus by immunofluorescent analysis.

Sample: Mock and treated HCT116 cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: p53 stained by p53 antibody [DO1] (GTX70214) diluted at 1:500.

Blue: Fluoroshield with DAPI (GTX30920).



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 27 Page 2 of 2