p53 antibody [DO1]

Cat. No. GTX70214

| Host | Mouse | References (40) |
|--------------|---|------------------------|
| Clonality | Monoclonal | 🚖 🚖 🚖 🊖 🪖 Review (2) |
| lsotype | lgG2a | Package |
| Applications | WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, ELISA | 100 μl |
| Reactivity | Human, Mouse | |

PRODUCT

Summary

p53 antibody recognizes p53 protein, which has a predicted molecular weight of 44 kDa but is typically detected by western blot at 53 kDa. p53, known widely as a tumor suppressor, plays critical roles in various cellular processes including cell cycle regulation, DNA repair, and apoptosis. p53 is mutated in many human cancers.

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500-1:10000 |
| ICC/IF | Assay dependent |
| IHC-P | 1:1000 |
| IHC-Fr | Assay dependent |
| FCM | Assay dependent |
| IP | 1:100-1:500 |
| ELISA | Assay dependent |

Note : Antigen retrieval can be done with EDTA at pH8.0 - Some customers have stated that the product works well without antigen retrieval.

Not tested in other applications.

| Calculated MW | 44 kDa. (<u>Note</u>) |
|---------------|-------------------------|
|---------------|-------------------------|

| iquid | |
|--|--|
| PBS, 20% Glycerol | |
| No Preservative | |
| Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. | |
| 0.25 mg/ml (Please refer to the vial label for the specific concentration.) | |
| | |




For full product information, images and publications, please visit our <u>website</u>.




| Immunogen | Recombinant human p53 expressed in E. coli. | |
|--------------|---|--|
| Purification | Protein A purified From tissue culture supernatant | |
| Conjugation | Unconjugated | |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. | |
| Note | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. | |

DATA IMAGES




GTX70214 WB Image

Non-transfected (–) and transfected (+) HCT116 whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with p53 antibody [DO1] (GTX70214) diluted at 1:1000. The HRPconjugated anti-mouse IgG antibody (GTX213111-01) was used to detect the primary antibody.



GTX70214 IHC-P Image

p53 antibody [DO1] detects p53 protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded human pancreatic cancer. p53 stained by p53 antibody [DO1] (GTX70214) diluted at 1:100. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX70214 IHC-P Image

p53 antibody [DO1] detects p53 protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded human NCI-H1299 xenograft. p53 stained by p53 antibody [DO1] (GTX70214) diluted at 1:100. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our <u>website</u>.