

CIDEc antibody

Cat. No. GTX70476

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human

Package
100 µg

Applications

Application Note

Western blot - Use at 1:500-1,000 dilution. However, the investigator should determine the optimal dilution for a specific application such as immunohistochemistry.

Calculated MW 27 kDa. ([Note](#))

Product Note Reacts with residues 10-25 [LLYPKSLSRHVSVRTS] of the 27 kDa human CIDEc protein. Cross reactivity to FSP27, which is the mouse homolog of CIDE-3: 93% sequence identity.

Properties

Form Liquid

Buffer PBS

Preservative 0.02% Sodium azide

Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Concentration 1 mg/ml (Please refer to the vial label for the specific concentration.)

Purification Affinity purified IgG

Conjugation Unconjugated

Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).