

CD19 antibody [HD37]

Cat. No. GTx72026

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-Fr, FCM
Reactivity	Human

Package
500 µl

Applications

Application Note

For a three-step avidin-biotin complex system, a dilution of 1:50 to 1:100 may be used as a guideline. Dilute according to the particular application being used. In general, the 0.05M Borate pH 8.0 containing 0.15M Sodium Chloride, 0.05% Sodium Azide, is a good diluent to use with most antibodies. Optimal dilutions/concentrations should be determined by the end user.

Properties

Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	0.17 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Hairy Cell Leukemia Cells
Purification	Protein A affinity purified
Conjugation	Unconjugated

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 12 Page 1 of 1