

## CD16 antibody [LNK16] (FITC)

## Cat. No. GTX74719

|              |                |
|--------------|----------------|
| Host         | Mouse          |
| Clonality    | Monoclonal     |
| Isotype      | IgG1           |
| Applications | FCM            |
| Reactivity   | Human, Primate |

References ( 1 )

Package

100 test

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution   |
|--------------------|--|
| FCM                | 20 µl reagent / 100 µl of whole blood or 10 <sup>6</sup> cells in a suspension |

Not tested in other applications.

## Properties


|   |   |
|---|---|
| Form  | Liquid  |
| Buffer  | PBS   |
| Preservative  | 15mM Sodium azide   |
| Storage   | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Protect from light.  |
| Concentration   | Batch dependent (Please refer to the vial label for the specific concentration.)  |
| Immunogen   | Normal human peripheral blood granulocytes  |
| Purification  | Purified by size-exclusion chromatography   |
| Conjugation   | Fluorescein isothiocyanate (FITC) <a href="#">Wavelength</a>  |
| For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |   |
| Note  | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 29 Page 1 of 2


## DATA IMAGES



## GTX74719 FCM Image

FACS analysis of human lymphocytes using GTX74719 CD16 antibody [LNK16] (FITC).


Antibody amount : 20  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood



## GTX74719 FCM Image

FACS analysis of human peripheral blood using GTX74719 CD16 antibody [LNK16] (FITC).

Antibody amount : 20  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood



## GTX74719 FCM Image

FACS analysis of human CD3 negative CD16 positive Lymphocytes (red-filled) and human CD3 positive CD16 negative Lymphocytes (black-dashed) using GTX74719 CD16 antibody [LNK16] (FITC).

Antibody amount : 20  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 29 Page 2 of 2